After Your Intravitreal Injection

What to Expect

Mild eye irritation/burning, frequent tearing, redness and transient floaters/bubbles can occur after an injection. These symptoms are usually mild and temporary, but can occasionally be severe. Most cases will improve overnight without additional treatment.

How to Care for Your Eye



Use artificial tears as needed for irritation. Rest the eye closed for 4–5 hours.



Do not rub or press on the treated eye.

Common and Temporary Symptoms



Redness
Burning or irritation is common, and often related to the antiseptic (Betadine) used to clean the eye.



Tearing
The eye may tear for a few hours after the injection.



Floaters or Blurry Vision
These symptoms usually improve by the next day.

When to Contact Us

Please call our office if you experience:



Severe pain and/or light sensitivity
that does not improve overnight.



New or worsening flashes/floaters
occurring or persisting >24 hours after the injection.



Significant vision loss.

HOUSTON RETINA ASSOCIATES

(281) 495-2222
www.hretina.com

Reviewed by
Henry Lin, MD, PhD

玻璃體腔注射後注意事項

注射後可能出現的症狀

注射後可能出現眼睛刺痛或灼熱、頻繁流淚、眼紅，以及短暫的飛蚊或氣泡。這些症狀通常輕微且短暫，但有時也可能較嚴重。大多數情況不需額外治療，隔夜便會改善。

如何照顧注射後的眼睛



眼睛不適時可使用人工淚液。注射後讓眼睛閉上休息 4–5 小時。



請勿揉擦或按壓接受注射的眼睛。

常見的暫時性症狀



眼紅

灼熱或刺激感很常見，通常與清潔眼睛所用的消毒液（Betadine）有關。



流淚

注射後眼睛可能會流淚數小時。



飛蚊或視力模糊

這些症狀通常會在隔天改善。

何時需要聯絡我們

若出現以下情況，請致電本診所：



嚴重眼痛和／或畏光

隔夜仍未改善。



新出現或加重的閃光、飛蚊

注射後超過 24 小時才出現，或持續超過 24 小時。



視力明顯下降

HOUSTON RETINA ASSOCIATES

(281) 495-2222

www.hretina.com

審閱：林醫師
Henry Lin, MD, PhD