After Your Intravitreal Injection

What to Expect

Mild eye irritation/burning, frequent tearing, redness and transient floaters/bubbles can occur after an injection. These symptoms are usually mild and temporary, but can occasionally be severe. Most cases will improve overnight without additional treatment.

How to Care for Your Eye



Use artificial tears as needed for irritation. Rest the eye closed for 4–5 hours.



Do not rub or press on the treated eye.

Common and Temporary Symptoms



Redness
Burning or irritation is common, and often related to the antiseptic (Betadine) used to clean the eye.



Tearing
The eye may tear for a few hours after the injection.



Floaters or Blurry Vision
These symptoms usually improve by the next day.

When to Contact Us

Please call our office if you experience:



Severe pain and/or light sensitivity
that does not improve overnight.



New or worsening flashes/floaters
occurring or persisting >24 hours after the injection.



Significant vision loss.

HOUSTON RETINA ASSOCIATES

(281) 495-2222
www.hretina.com

Reviewed by
Henry Lin, MD, PhD

Después de su Inyección Intravítrea

Qué Puede Esperar

Es común tener irritación o ardor leve en el ojo, lagrimeo frecuente, enrojecimiento y aparición temporal de moscas volantes o burbujas después de una inyección. Estos síntomas suelen ser leves y temporales, pero en ocasiones pueden ser más intensos. La mayoría de los casos mejorarán durante la noche sin necesidad de tratamiento adicional.

Cómo Cuidar su Ojo



Use lágrimas artificiales según sea necesario para la irritación. Mantenga el ojo cerrado durante 4 a 5 horas.



No se frote ni ejerza presión sobre el ojo tratado.

Síntomas Comunes y Temporales



Enrojecimiento
Es común tener ardor o irritación y a menudo se debe al antiséptico (Betadine) utilizado para limpiar el ojo.



Lagrimeo
El ojo puede llorar durante algunas horas después de la inyección.



Moscas volantes o visión borrosa
Estos síntomas suelen mejorar para el día siguiente.

Cuándo Comunicarse con Nosotros

Llame a nuestra oficina si presenta:



Dolor intenso y/o sensibilidad a la luz
que no mejora durante la noche.



Aparición o empeoramiento de destellos o moscas volantes
que persisten por más de 24 horas después de la inyección.



Pérdida significativa de la visión.

HOUSTON RETINA ASSOCIATES

(281) 495-2222
www.hretina.com

Revisado por
Henry Lin, MD, PhD